

HIV Protocol: For HIV ELISA and Western Blot Positive Patients
HIV-infected 住院病人 (Baseline) 98.01.05 修訂

Test	Remark
CBC/DC	
Lymphocyte markers, infection (請用核醫碼開立 88060E)	* 2 nd test is recommended <u>PRIOR</u> to initiation of HAART or * 2 nd test is recommended 1 month after initial test
HIV viral load (主治醫師的碼才能開)	* 2 nd test is recommended <u>PRIOR</u> to initiation of HAART or * 2 nd test is recommended 1 month after initial test
Chest X-ray (PA and lateral view)	
VDRL	若陽性，加測 TPHA
**Anti-HAV	Vaccinate if IgG negative (OPD)
**HBsAg	若是陰性，加測 anti-HBsAb and anti-HBc IgG Ab 或 anti-HBc IgM Ab (acute HBV infection) 若是陽性，加測 HBeAg and HDV Ab
Anti-HCV Ab	HCV RNA (if acute HCV infection is suspected, 需自費)
Serum Cryptococcal antigen	
Toxoplasma IgG Ab	IgM testing depending on clinical status
CMV IgG Ab	IgM testing depending on clinical status
Serum chemistry 包括 ALT, AST, LDH, BUN, Crea, Na, K, T-bil, ALP, Glu, lipid profile	Repeat lipid profile and glucose prior to initiation of HAART
Pap smear + Vaginal secretion for Trichomonas	For women only
Urinalysis	
Ophthalmology referral	If CD4<50
Urine CMV	If CD4<50
Stool routine	
Serum amoeba IHA	
Sputum TB culture X 3 sets	
EKG	

** Vaccinate if negative

