

高雄榮民總醫院

口咽癌診療原則

2018年01月03日第一版

頭頸癌醫療團隊擬訂

注意事項：這個診療原則主要作為醫師和其他保健專家診療癌症病人參考之用。假如你是一個癌症病人，直接引用這個診療原則並不恰當，只有你的醫師才能決定給你最恰當的治療。

會議討論

上次會議：2017/05/17

本共識與上一版的差異

上一版	新版
<ol style="list-style-type: none">1. 未公告AJCC 8th2. 未以p16病理染色決定分期。3. Adverse feature：ENE(+)為N3b。4. 無區分P16。5. 未標示M1之治療指引及第一線C/T	<ol style="list-style-type: none">1. 全面改以AJCC第8版為分期依據2. 將P16檢查結果納入分期決定項目3. 移除Adverse feature中ENE(+)，直接改為N3b（會議過程保留此項）4. 若P16(+)則移除T4b/N2abc/N3ab5. 新增M1之治療指引及第一線C/T建議

Oropharyngeal (P16 negative) cancer Clinical staging AJCC 8th

Oropharyngeal (p16 negative) cancer TNM clinical staging AJCC UICC 2017

Primary tumor (T)	
Oropharynx (p16-)	
T category	T criteria
TX	Primary tumor cannot be assessed
Tis	Carcinoma <i>in situ</i>
T1	Tumor 2 cm or smaller in greatest dimension
T2	Tumor larger than 2 cm but not larger than 4 cm in greatest dimension
T3	Tumor larger than 4 cm in greatest dimension or extension to lingual surface of epiglottis
T4	Moderately advanced or very advanced local disease
T4a	Moderately advanced local disease. Tumor invades the larynx, extrinsic muscle of tongue, medial pterygoid, hard palate, or mandible.*
T4b	Very advanced local disease. Tumor invades lateral pterygoid muscle, pterygoid plates, lateral nasopharynx, or skull base or encases carotid artery.
* NOTE: Mucosal extension to lingual surface of epiglottis from primary tumors of the base of the tongue and vallecula does not constitute invasion of the larynx.	
Regional lymph nodes (N)	
Clinical N (cN) - Oropharynx (p16-) and hypopharynx	
N category	N criteria
NX	Regional lymph nodes cannot be assessed
N0	No regional lymph node metastasis
N1	Metastasis in a single ipsilateral lymph node, 3 cm or smaller in greatest dimension and ENE(-)
N2	Metastasis in a single ipsilateral node larger than 3 cm but not larger than 6 cm in greatest dimension and ENE(-); or metastases in multiple ipsilateral lymph nodes, none larger than 6 cm in greatest dimension and ENE(-); or in bilateral or contralateral lymph nodes, none larger than 6 cm in greatest dimension and ENE(-)
N2a	Metastasis in a single ipsilateral node larger than 3 cm but not larger than 6 cm in greatest dimension and ENE(-)
N2b	Metastasis in multiple ipsilateral nodes, none larger than 6 cm in greatest dimension and ENE(-)
N2c	Metastasis in bilateral or contralateral lymph nodes, none larger than 6 cm in greatest dimension and ENE(-)
N3	Metastasis in a lymph node larger than 6 cm in greatest dimension and ENE(-); or metastasis in any node(s) and clinically overt ENE(+)
N3a	Metastasis in a lymph node larger than 6 cm in greatest dimension and ENE(-)
N3b	Metastasis in any node(s) and clinically overt ENE(+)
NOTE: A designation of "U" or "L" may be used for any N category to indicate metastasis above the lower border of the cricoid (U) or below the lower border of the cricoid (L). Similarly, clinical and pathological ENE should be recorded as ENE(-) or ENE(+).	

Distant metastasis (M)			
Oropharynx (p16-) and hypopharynx			
M category	M criteria		
M0	No distant metastasis		
M1	Distant metastasis		
Prognostic stage groups			
When T is...	And N is...	And M is...	Then the stage group is...
Tis	N0	M0	0
T1	N0	M0	I
T2	N0	M0	II
T3	N0	M0	III
T1, T2, T3	N1	M0	III
T4a	N0, 1	M0	IVA
T1, T2, T3, T4a	N2	M0	IVA
Any T	N3	M0	IVB
T4b	Any N	M0	IVB
Any T	Any N	M1	IVC

TNM: tumor, node, metastasis; AJCC: American Joint Committee on Cancer; UICC: Union for International Cancer Control; ENE: extranodal extension.

Oropharyngeal (P16 negative) cancer

Pathological staging AJCC 8th

Oropharyngeal (p16 negative) cancer TNM pathological staging AJCC UICC 2017

Primary tumor (T)	
Oropharynx (p16-)	
T category	T criteria
TX	Primary tumor cannot be assessed
Tis	Carcinoma <i>in situ</i>
T1	Tumor 2 cm or smaller in greatest dimension
T2	Tumor larger than 2 cm but not larger than 4 cm in greatest dimension
T3	Tumor larger than 4 cm in greatest dimension or extension to lingual surface of epiglottis
T4	Moderately advanced or very advanced local disease
T4a	Moderately advanced local disease. Tumor invades the larynx, extrinsic muscle of tongue, medial pterygoid, hard palate, or mandible.*
T4b	Very advanced local disease. Tumor invades lateral pterygoid muscle, pterygoid plates, lateral nasopharynx, or skull base or encases carotid artery.
* NOTE: Mucosal extension to lingual surface of epiglottis from primary tumors of the base of the tongue and vallecula does not constitute invasion of the larynx.	
Regional lymph nodes (N)	
Pathological N (pN) - Oropharynx (p16-) and hypopharynx	
N category	N criteria
NX	Regional lymph nodes cannot be assessed
N0	No regional lymph node metastasis
N1	Metastasis in a single ipsilateral lymph node, 3 cm or smaller in greatest dimension and ENE(-)
N2	Metastasis in a single ipsilateral lymph node, 3 cm or smaller in greatest dimension and ENE(+); or larger than 3 cm but not larger than 6 cm in greatest dimension and ENE(-); or metastases in multiple ipsilateral lymph nodes, none larger than 6 cm in greatest dimension and ENE(-); or in bilateral or contralateral lymph node(s), none larger than 6 cm in greatest dimension and ENE(-)
N2a	Metastasis in single ipsilateral node 3 cm or smaller in greatest dimension and ENE(+); or a single ipsilateral node larger than 3 cm but not larger than 6 cm in greatest dimension and ENE(-)
N2b	Metastasis in multiple ipsilateral nodes, none larger than 6 cm in greatest dimension and ENE(-)
N2c	Metastasis in bilateral or contralateral lymph node(s), none larger than 6 cm in greatest dimension and ENE(-)
N3	Metastasis in a lymph node larger than 6 cm in greatest dimension and ENE(-); or in a single ipsilateral node larger than 3 cm in greatest dimension and ENE(+); or multiple ipsilateral, contralateral or bilateral nodes, any with ENE(+); or a single contralateral node 3 cm or smaller and ENE(+)
N3a	Metastasis in a lymph node larger than 6 cm in greatest dimension and ENE(-)
N3b	Metastasis in a single ipsilateral node larger than 3 cm in greatest dimension

	and ENE(+); or multiple ipsilateral, contralateral or bilateral nodes, any with ENE(+); or a single contralateral node 3 cm or smaller and ENE(+)		
NOTE: A designation of "U" or "L" may be used for any N category to indicate metastasis above the lower border of the cricoid (U) or below the lower border of the cricoid (L). Similarly, clinical and pathological ENE should be recorded as ENE(-) or ENE(+).			
Distant metastasis (M)			
Oropharynx (p16-) and hypopharynx			
M category	M criteria		
M0	No distant metastasis		
M1	Distant metastasis		
Prognostic stage groups			
When T is...	And N is...	And M is...	Then the stage group is...
Tis	N0	M0	0
T1	N0	M0	I
T2	N0	M0	II
T3	N0	M0	III
T1, T2, T3	N1	M0	III
T4a	N0, 1	M0	IVA
T1, T2, T3, T4a	N2	M0	IVA
Any T	N3	M0	IVB
T4b	Any N	M0	IVB
Any T	Any N	M1	IVC

TNM: tumor, node, metastasis; AJCC: American Joint Committee on Cancer; UICC: Union for International Cancer Control; ENE: extranodal extension.

Oropharyngeal (P16 Positive) cancer

Pathological staging AJCC 8th

HPV related oropharyngeal carcinoma TNM clinical staging AJCC UICC 2017

Primary tumor (T)			
T category	T criteria		
T0	No primary identified		
T1	Tumor 2 cm or smaller in greatest dimension		
T2	Tumor larger than 2 cm but not larger than 4 cm in greatest dimension		
T3	Tumor larger than 4 cm in greatest dimension or extension to lingual surface of epiglottis		
T4	Moderately advanced local disease. Tumor invades the larynx, extrinsic muscle of tongue, medial pterygoid, hard palate, or mandible or beyond.*		
Regional lymph nodes (N) - Clinical N (cN)			
N category	N criteria		
NX	Regional lymph nodes cannot be assessed		
N0	No regional lymph node metastasis		
N1	One or more ipsilateral lymph nodes, none larger than 6 cm		
N2	Contralateral or bilateral lymph nodes, none larger than 6 cm		
N3	Lymph node(s) larger than 6 cm		
Distant metastasis (M)			
M category	M criteria		
M0	No distant metastasis		
M1	Distant metastasis		
Prognostic stage groups - Clinical			
When T is...	And N is...	And M is...	Then the stage group is...
T0, T1, or T2	N0 or N1	M0	I
T0, T1, or T2	N2	M0	II
T3	N0, N1, or N2	M0	II
T0, T1, T2, T3, or T4	N3	M0	III
T4	N0, N1, N2, or N3	M0	III
Any T	Any N	M1	IV

HPV: human papillomavirus; TNM: tumor, node, metastasis; AJCC: American Joint Committee on Cancer; UICC: Union for International Cancer Control.

* Mucosal extension to lingual surface of epiglottis from primary tumors of the base of the tongue and vallecula does not constitute invasion of the larynx.

HPV related oropharyngeal carcinoma TNM pathologic staging AJCC UICC 2017

Primary tumor (T)			
T category	T criteria		
T0	No primary identified		
T1	Tumor 2 cm or smaller in greatest dimension		
T2	Tumor larger than 2 cm but not larger than 4 cm in greatest dimension		
T3	Tumor larger than 4 cm in greatest dimension or extension to lingual surface of epiglottis		
T4	Moderately advanced local disease. Tumor invades the larynx, extrinsic muscle of tongue, medial pterygoid, hard palate, or mandible or beyond.*		
Regional lymph nodes (N) - Pathological N (pN)			
N category	N criteria		
NX	Regional lymph nodes cannot be assessed		
pN0	No regional lymph node metastasis		
pN1	Metastasis in four or fewer lymph nodes		
pN2	Metastasis in more than four lymph nodes		
Distant metastasis (M)			
M category	M criteria		
M0	No distant metastasis		
M1	Distant metastasis		
Prognostic stage groups - Pathological			
When T is...	And N is...	And M is...	Then the stage group is...
T0, T1, or T2	N0, N1	M0	I
T0, T1, or T2	N2	M0	II
T3 or T4	N0, N1	M0	II
T3 or T4	N2	M0	III
Any T	Any N	M1	IV

HPV: human papillomavirus; TNM: tumor, node, metastasis; AJCC: American Joint Committee on Cancer; UICC: Union for International Cancer Control.

* Mucosal extension to lingual surface of epiglottis from primary tumors of the base of the tongue and vallecula does not constitute invasion of the larynx.


Carcinoma of Oropharynx

高雄榮民總醫院 臨床診療指引 Ver.1 修訂於 2018.01.03 Page 1 (Ref. 1,2)

WORK-UP	STAGING & TREATMENT	FOLLOW-UP
<ul style="list-style-type: none">• <u>History & PE</u>• <u>Biopsy & Pathology</u>• <u>Image</u><ul style="list-style-type: none">→ MRI or CT of H & N→ Chest X-ray→ Bone scan→ Abd. Sono→ ± Neck Sono→ ± PET scan• <u>Dental evaluation</u><ul style="list-style-type: none">→ Panorex→ ± teeth extraction• UGI series exam• <u>Multidisciplinary consultation</u><ul style="list-style-type: none">± Swallowing evaluation	<ul style="list-style-type: none">• <u>[T1-2, N0-1, M0]</u> 詳見 <i>Page 2</i>• <u>[T3-4a, N0-1, M0]</u> 詳見 <i>Page 3</i>• <u>[Any T, N2-3, M0]</u> 詳見 <i>Page 4</i>• <u>Very advanced stage</u> 詳見 <i>Page 5</i>	<ul style="list-style-type: none">• <u>[Post-Tx within 6 months]</u><ul style="list-style-type: none">→ Every 1-2 months: PE→ Baseline MRI or CT• <u>[0.5-3 years after Tx]</u><ul style="list-style-type: none">→ Every 2-3 months: PE→ Every 1 year: H & N MRI or CT, CxR, Bone scan & Abd. Sono ± Neck Sono as clinically indicated• <u>[3-5 years after Tx]</u><ul style="list-style-type: none">→ Every 4-6 months: PE• <u>[5 years later after Tx]</u><ul style="list-style-type: none">→ Every 6-12 months: PE

Carcinoma of Oropharynx

高雄榮民總醫院 臨床診療指引 Ver.1 修訂於 2018.01.03 Page 2 (Ref. 1,2)




#可考慮Elective neck dissection或close follow-up

*Adverse features: Extranodal extension, positive or close margins, N2 or N3 nodal disease, nodal disease in levels IV or V, perineural invasion, lymphovascular invasion.

Carcinoma of Oropharynx

高雄榮民總醫院 臨床診療指引 Ver.1 修訂於 2018.01.03 Page 3 (Ref. 1,2)




#Neck dissection level 依cN status而定。

*Adverse features : Extranodal extension, positive margins, N2 or N3 nodal disease, nodal disease in levels IV or V, perineural invasion, lymphovascular invasion.

Carcinoma of Oropharynx

高雄榮民總醫院 臨床診療指引 Ver.1 修訂於 2018.01.03 Page 4 (Ref. 1,2)




Neck dissection level 依primary部位及cN status而定。

* Adverse features : Extranodal extension, positive margins, pT3-4, N2-3 nodal disease, nodal disease in levels IV or V, perineural invasion, lymphovascular invasion.

Carcinoma of Oropharynx

高雄榮民總醫院 臨床診療指引 Ver.1 修訂於 2018.01.03 Page 6 (Ref. 1,2)




ECOG Performance Status 0-1 註6

* ECOG Performance Status 2

\$ ECOG Performance Status 3

Carcinoma of Oropharynx

高雄榮民總醫院 臨床診療指引 Ver.1 修訂於 2018.01.03 Page 7 (Ref. 1,15-17)



ECOG Performance Status 0-1 註6

* ECOG Performance Status 2

\$ ECOG Performance Status 3

Carcinoma of Oropharynx

高雄榮民總醫院 臨床診療指引 Ver.1 修訂於 2018.01.03 Page 8 (Ref. 3-5)

註1

Principles of Radiotherapy

Definitive Radiotherapy

- Primary and gross adenopathy : 66 - 74 Gy (1.8-2.0 Gy/fraction)
- Neck uninvolved nodal stations : 44 - 64 Gy (1.6-2.0 Gy/fractions)

Postoperative Radiotherapy

- Preferred interval between operation and radiotherapy is ≤ 6 weeks.
- Primary : 60-66 Gy (1.8-2.0 Gy/fraction)
- Neck involved nodal stations : 60 - 66 Gy (1.8-2.0 Gy/fraction)
- Neck uninvolved nodal stations : 44 - 64 Gy (1.6-2.0 Gy/fraction)

CCRT or RT

- RT alone if : old age, impaired renal function, poor condition or refused chemotherapy

Palliative RT

- Indicated in : relieve local symptoms, prevent debilitation such as spinal cord compression and pathological fracture, achieve durable locoregional control.

Carcinoma of Oropharynx

註2

高雄榮民總醫院 臨床診療指引 Ver.1 修訂於 2018.01.03 Page 9 (Ref. 6-10)

Principles of Chemotherapy

Concurrent with RT

Regimen 1: q3w CDDP ± Cetuximab^{註5} + RT

- Cisplatin (80-100mg/ m²) q3w during R/T
- Cetuximab(400mg/ m²) loading dose first week, then Cetuximab(250mg/ m²) maintain dose D1 + Cisplatin (80-100mg/ m²) q3w D2 during R/T

Regimen 2: Weekly CDDP ± Cetuximab^{註5} + RT

- Cisplatin (30-40mg/ m²) weekly during R/T
- Cetuximab(400mg/ m²) loading dose first week, and then Cisplatin (30-40mg/ m²) weekly D1 + Cetuximab(250mg/ m²) maintain dose D2 during R/T

Regimen 3: q3w Carboplatin^{註4} ± Cetuximab^{註5} + RT

- Carboplatin (AUC x 5mg) q3w during R/T
- Cetuximab(400mg/ m²) loading dose first week, then Cetuximab(250mg/ m²) maintain dose D1 + Carboplatin (AUC x 5mg) q3w D2 during R/T

Regimen 4: Weekly Cetuximab^{註5} + RT

- Cetuximab(400mg/ m²) loading dose first week, then Cetuximab(250mg/ m²) maintain dose during RT

Carcinoma of Oropharynx

註3

高雄榮民總醫院 臨床診療指引 Ver.1 修訂於 2018.01.03 Page 10 (Ref. 11-15)

Regimens of Chemotherapy

Induction, adjuvant, 建議2-3cycles

Regimen 1: q3-4 weeks CDDP ± F ± weekly Cetuximab^{註5}

- Cisplatin(80-100mg/ m²) D1
- Fluorouracil (5-FU) (600-1000 mg/m²) D2-D5
- Cetuximab(400mg/ m²) loading dose first week, then weekly Cetuximab (250mg/ m²)

Regimen 2: q3-4 weeks P ± F ± weekly Cetuximab^{註5}

- Cisplatin (20mg/ m²) D1-D5
- Fluorouracil (5-FU) (600-1000 mg/ m²) D1-D5
- Cetuximab(400mg/ m²) loading dose first week, then weekly Cetuximab (250mg/ m²)

Regimen 3: q3-4 weeks Carboplatin ± F ± weekly Cetuximab^{註5}

- Carboplatin (AUC x 5mg) D1
- Fluorouracil (5-FU) (600-1000 mg/ m²) D2-D5
- Cetuximab(400mg/ m²) loading dose first week, then weekly Cetuximab (250mg/ m²)

Carcinoma of Oropharynx

註3

高雄榮民總醫院 臨床診療指引 Ver.1 修訂於 2018.01.03 Page 11 (Ref. 11-15)

Regimens of Chemotherapy

Induction, adjuvant, 建議2-3cycles

Regimen 4: q3-4 weeks T + P ± F ± weekly Cetuximab^{註5}

- Taxotere(60 mg/ m²) D1
- Cisplatin(60 mg/ m²) D1
- Fluorouracil (5-FU) (600 mg/m²) D1-D5
- Cetuximab (400mg/ m²) loading dose first week, then Cetuximab (250mg/ m²) maintain dose

Regimen 5: weekly Cetuximab^{註5}

- Cetuximab (400mg/ m²) loading dose first week, then Cetuximab (250mg/ m²) maintain dose

Regimen 6: oral Fluorouracil

- Ufur cap (tegafur 100mg+uracil 224mg) 2# BID-TID
(Salvage or palliative CT中作為取代iv-formed 5-FU之替代藥物)

Regimen 7: weekly Methotrexate

- Methotrexate (40-60mg/ m²)

Carcinoma of Oropharynx

註4

高雄榮民總醫院 臨床診療指引 Ver.1 修訂於 2018.01.03 Page 12 (Ref. 16,17)

Regimens of Chemotherapy

Recurrent, unresectable, metastatic

Regimen 1: q3-4 weeks CDDP ± F ± weekly Cetuximab^{註5}

- Cisplatin(80-100mg/ m²) D1 or Cisplatin (20mg/ m²) D1-D5
- Fluorouracil (5-FU) (600-1000 mg/m²) D2-D5
- Cetuximab(400mg/ m²) loading dose first week, then weekly Cetuximab (250mg/ m²)

Regimen 2: q3-4 weeks Carboplatin^{註6} ± F ± weekly Cetuximab^{註5}

- Carboplatin (AUC x 5mg) D1
- Fluorouracil (5-FU) (1000mg/ m²) D2-D5
- Cetuximab(400mg/ m²) loading dose first week, then weekly Cetuximab (250mg/ m²)

Regimen 3: q3-4 weeks T^{註5} ± P

- Taxotere(60 mg/ m²) D1
- Cisplatin(60-75 mg/ m²) D1

Regimen 4: q3-4 weeks T^{註5} ± Carboplatin

- Taxotere(60 mg/ m²) D1
- Carboplatin (AUC x 5mg) D1

Carcinoma of Oropharynx

註5

高雄榮民總醫院 臨床診療指引 Ver.1 修訂於 2018.01.03 Page 13

特殊用藥健保給付規定

Taxotere

- 頭頸部癌，限局部晚期且無遠端轉移之頭頸部鱗狀細胞癌且無法手術切除者。
- 與Cisplatin 及5-FU 併用，作為放射治療前的引導治療，限使用四個療程。

Cetuximab

- 限與放射線療法合併使用於局部晚期之口咽癌、下咽癌及喉癌患者，且符合下列條件之一：
 1. 年齡 ≥ 70 歲
 2. $Ccr < 50ml/min$
 3. 聽力障礙者 (聽力障礙定義為500Hz、1000Hz、2000Hz 平均聽力損失大於25 分貝)
 4. 無法耐受platinum-based 化學治療。
- 使用總療程以接受8 次輸注為上限。
- 需經事前審查核准後使用。

Carboplatin

- 限腎功能不佳 ($CCr < 60$) 或曾作單側或以上腎切除之惡性腫瘤患者使用。

Carcinoma of Oropharynx

註6

高雄榮民總醫院 臨床診療指引 Ver.1 修訂於 2018.01.03 Page 14

Eastern Cooperative Oncology Group (ECOG) Performance Status

Grade	Description	Suggestion
0	Normal activity fully ambulatory (無症狀)	按照標準化療評估及療程。
1	Symptoms, but nearly fully ambulatory (有症狀，完全步行，但對生活無影響)	按照標準化療評估及療程。
2	Some bed time, but needs to be in bed less than 50% of normal daytime (躺在床上的時間<50%)	按照標準化療評估及療程。
3	Needs to be in bed more than 50% of normal daytime (躺在床上的時間>50%)	可視情況考慮停止化學治療。
4	Unable to get out of bed (長期完全臥床)	建議停止化學治療。
5	Dead	

Carcinoma of Oropharynx

References

高雄榮民總醫院 臨床診療指引 Ver.1 修訂於 2018.01.03 Page 15

1. NCCN Clinical Practice Guidelines in Oncology – Head and Neck Cancers Version 2. 2017
2. AJCC (American Joint Committee on Cancer) Manual for Staging of Cancer, 8th ed, Amin M, Edge S, Greene F, et al. (Eds), Springer-Verlag, New York 2017.
3. Peter AP, Michael EC, Greg D, et al. Up-front neck dissection followed by concurrent chemoradiation in patients with regionally advanced head and neck cancer. *Head Neck*. 2012;34:1798-1803
4. Bradley, PJ, MacLennan, K, Brakenhoff, RH, Leemans, CR. Status of primary tumour surgical margins in squamous head and neck cancer: prognostic implications. *Curr Opin Otolaryngol Head Neck Surg* 2007; 15:74.
5. Vermorken JB, Remenar E, van Herpen C, Gorlia T, Mesia R, Degardin M, Stewart JS, Jelic S, Betka J, Preiss JH, et al. Cisplatin, fluorouracil, and docetaxel in unresectable head and neck cancer. *N Engl J Med*. 2007 Oct 25; 357(17):1695-704
6. Forastiere AA, Zhang Q, Weber RS, et al. Long-term results of RTOG 91-11: a comparison of three nonsurgical treatment strategies to preserve the larynx in patients with locally advanced larynx cancer. *J Clin Oncol* 2013;31:845-852.
7. Adelstein DJ, Li Y, Adams GL, et al. An intergroup phase III comparison of standard radiation therapy and two schedules of concurrent chemoradiotherapy in patients with unresectable squamous cell head and neck cancer. *J Clin Oncol* 2003;21(1):92-98.
8. Bonner JA, Harari PM, Giralt J, et al. Radiotherapy plus cetuximab for locoregionally advanced head and neck cancer: 5-year survival data from a phase 3 randomised trial, and relation between cetuximab-induced rash and survival. *Lancet Oncol* 2010;11:21-28
9. Denis F, Garaud P, Bardet E, et al. Final results of the 94-01 French Head and Neck Oncology and Radiotherapy Group randomized trial comparing radiotherapy alone with concomitant radiochemotherapy in advanced-stage oropharynx carcinoma. *J Clin Oncol* 2004;22:69-76
10. Bourhis J, Sire C, Graff P, et al. Concomitant chemoradiotherapy versus acceleration of radiotherapy with or without concomitant chemotherapy in locally advanced head and neck carcinoma (GORTEC 99-02): an open-label phase 3 randomised trial. *Lancet Oncol* 2012;13:145-153.
11. Forastiere AA, Zhang Q, Weber RS, et al. Long-term results of RTOG 91-11: a comparison of three nonsurgical treatment strategies to preserve the larynx in patients with locally advanced larynx cancer. *J Clin Oncol* 2013;31:845-852.
12. Adelstein DJ, Li Y, Adams GL, et al. An intergroup phase III comparison of standard radiation therapy and two schedules of concurrent chemoradiotherapy in patients with unresectable squamous cell head and neck cancer. *J Clin Oncol* 2003;21(1):92-98.
13. Bonner JA, Harari PM, Giralt J, et al. Radiotherapy plus cetuximab for locoregionally advanced head and neck cancer: 5-year survival data from a phase 3 randomised trial, and relation between cetuximab-induced rash and survival. *Lancet Oncol* 2010;11:21-28. Denis F, Garaud P, Bardet E, et al. Final results of the 94-01 French Head and Neck Oncology and Radiotherapy Group randomized trial comparing radiotherapy alone with concomitant radiochemotherapy in advanced-stage oropharynx carcinoma. *J Clin Oncol* 2004;22:69-76.
14. Bourhis J, Sire C, Graff P, et al. Concomitant chemoradiotherapy versus acceleration of radiotherapy with or without concomitant chemotherapy in locally advanced head and neck carcinoma (GORTEC 99-02): an open-label phase 3 randomised trial. *Lancet Oncol* 2012;13:145-153.
15. Posner MR, Hershock DM, Blajman CR, Mickiewicz E, Winkquist E, Gorbounova V, Tjulandin S, Shin DM, Cullen K, Ervin TJ, et al. Cisplatin and fluorouracil alone or with docetaxel in head and neck cancer. *N Engl J Med*. 2007 Oct 25; 357(17):1705-1
16. Vermorken JB, Mesia, R, Rivera F, Platinum-Based Chemotherapy plus Cetuximab in Head and Neck Cancer *N Engl J Med*. 2008 Sep 11; 359:1116-27
17. Guigay J, Fayette J, Dillies A-F, et al. Cetuximab, docetaxel, and cisplatin (TPEX) as first-line treatment in patients with recurrent or metastatic (R/M) squamous cell carcinoma of the head and neck (SCCHN): Final results of phase II trial GORTEC 2008-03 [abstract]. *J Clin Oncol* 2012;30(Suppl 15):Abstract 5505.