

高雄榮民總醫院

皮膚癌(BCC)診療原則

修訂日期:2024.04.19

癌委會公告日期:2024.06.03

BCC診療指引審視修訂會議討論日期


● 上次會議： 2023/02/12

上一版	新版
NCCN Guidelines 2023年版	更換附件為:NCCN Guidelines 2024年版



基底細胞癌(BCC)

診斷	初步評估	分期	初始治療	療效評估	輔助治療	追蹤
----	------	----	------	------	------	----



§ : Image studies is indicated for extensive disease (deep structural involvement such as bone, deep soft tissue, perineural disease)


+ : regional or distal metastatic disease(初始皮膚病灶治療同局限型)

☆ : 附件一

△ : 附件二

: T any, N0, M0(附件三)






BCC-1

- Preliminary Workup:
 - ▶ Bullet removed: Complete skin exam.
 - ▶ Bullet added: Shave excision if applicable.
- New option header added: Additional Workup
 - ▶ New option added: Complete skin examination.
- Risk Status, middle pathway following High risk, new option added: Consider imaging if clinical exam insufficient for following disease.
- Footnote d revised: Extensive disease includes deep structural involvement such as bone, named nerves perineural disease, and deep soft tissue. If perineural disease of named nerve(s) is suspected, MRI with contrast is preferred. If bone disease is suspected, CT with contrast is preferred unless contraindicated.





^b Principles of Pathology (BCC-A).

^c Risk Factors for Recurrence (BCC-B).

^h Principles of Treatment (BCC-C).

ⁱ In patients with superficial basal cell skin cancer, therapies such as topical imiquimod, topical 5-fluorouracil, photodynamic therapy, or cryotherapy may be considered, although cure rates are approximately 10% lower than for surgical treatment modalities. Jansen MHE, et al. J Invest Dermatol 2018;138:527-533. Drew BA, et al. Dermatol Surg 2017;43:1423-1430.

^j Principles of Radiation Therapy (BCC-D).

^k PDEMA with permanent section analysis or intraoperative frozen section analysis is an alternative to Mohs. See Principles of PDEMA Technique (SCC-G). Mohs surgery should be performed by dermatologic surgeons who have specialized training and experience in this procedure.

^m As per other appropriate use criteria. Task Force/Committee Members, Vidal CI, Armbract EA, Andrea AA, et al. J Am Acad Dermatol 2019;80:189-207.e11.

Note: All recommendations are category 2A unless otherwise indicated.
Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.

BCC-2

Primary Treatment:

- ▶ Top option revised: Curettage and electrodesiccation (C&E) or shave excision—Excluding terminal hair-bearing areas, such as the scalp, pubic and axillary regions, and beard area in males: or.
- ▶ Second option revised: If tumor appears to extend beyond the dermis, surgical or shave excision should generally be performed rather than C&E.
- ▶ New option added: Topical therapy.


Footnote removed: For tumors on cheeks, forehead, scalp, neck, and pretibia that are <6 mm in depth and confined to the dermis, C&E may be considered as an alternative primary treatment option if Mohs, resection with PDEMA, and standard excision are not feasible due to patient comorbidities. See Risk Factors for Recurrence (BCC-B).

New footnotes added:

- ▶ Footnote l: Mohs surgery should be performed by dermatologic surgeons who have specialized training and experience in this procedure. (Also page BCC-3A)
- ▶ Footnote m: As per other appropriate use criteria. Task Force/Committee Members, Vidal CL, Armbract EA, Andrea AA, et al. J Am Acad Dermatol 2019;80:189-207.e11. (Also page BCC-3A)



皮膚癌
多專科團隊



Footnotes on [BCC-3A](#)

Note: All recommendations are category 2A unless otherwise indicated.
Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.





STRATIFICATION TO DETERMINE TREATMENT OPTIONS FOR LOCAL BCC BASED ON RISK FACTORS FOR RECURRENCE^a

Risk Group	Low Risk	High Risk
Treatment options	BCC-2	BCC-3
H&P		
Location/size	Trunk, extremities <2 cm	Trunk, extremities ≥2 cm Head, neck, hands, feet, pretibia, and anogenital (any size) ^c
Borders	Well-defined	Poorly defined
Primary vs. recurrent	Primary	Recurrent
Immunosuppression	(-)	(+)
Site of prior RT	(-)	(+)
Pathology (BCC-A)		
Subtype	Nodular, superficial ^b	Aggressive growth pattern ^d
Perineural involvement	(-)	(+)

^a Any high-risk factor places the patient in the high-risk category.

^b Low-risk histologic subtypes include nodular, superficial, and other non-aggressive growth patterns such as keratotic, infundibulocystic, and fibroepithelioma of Pinkus.

^c This area constitutes high risk based on location, independent of size. Narrow excision margins due to anatomic and functional constraints are associated with increased recurrence rates with standard histologic processing. Complete margin assessment such as with Mohs or PDEMA is recommended for optimal tumor clearance and maximal tissue conservation. For tumors <6 mm in size, without other high-risk features, other treatment modalities may be considered if at least 4-mm clinically tumor-free margins can be obtained without significant anatomic or functional distortions.

^d Having basosquamous, infiltrative, sclerosing/morpheaform, micronodular, and BCC with carcinosarcomatous differentiation features in any portion of the tumor. In some cases, basosquamous tumors may be prognostically similar to squamous cell carcinoma (SCC); clinicopathologic correlation is recommended in these cases to further consider prognostic implication.

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



STRATIFICATION TO DETERMINE TREATMENT OPTIONS FOR LOCAL BCC BASED ON RISK FACTORS FOR RECURRENCE¹

Risk Group	Low Risk	High Risk
Treatment Options	See BCC-2	See BCC-3
H&P		
Location/size	Trunk, extremities <2 cm	Trunk, extremities ≥2 cm <u>Cheeks, forehead, scalp, neck, and pretibia (any size)</u> Head, neck, hands, feet, pretibia, and anogenital (any size) ³
Borders	Well-defined	Poorly defined
Primary vs. recurrent	Primary	Recurrent
Immunosuppression	(-)	(+)
Site of prior RT	(-)	(+)
Pathology (See BCC-A)		
Subtype	Nodular, superficial ²	Aggressive growth pattern ⁴
Perineural involvement	(-)	(+)

¹ Any high-risk factor places the patient in the high-risk category.

² Low-risk histologic subtypes include nodular, superficial, and other non-aggressive growth patterns such as keratotic, infundibulocystic, and fibroepithelioma of Pinkus.


³ This area constitutes high risk based on location, independent of size. Narrow excision margins due to anatomic and functional constraints are associated with increased recurrence rates with standard histologic processing. Complete margin assessment such as with Mohs or PDEMA is recommended for optimal tumor clearance and maximal tissue conservation. For tumors <6 mm in size, without other high-risk features, other treatment modalities may be considered if at least 4-mm clinically tumor-free margins can be obtained without significant anatomic or functional distortions.

⁴ Having (mixed) infiltrative, micronodular, morpheaform, basosquamous, sclerosing, or carcinosarcomatous differentiation features in any portion of the tumor. In some cases basosquamous tumors may be prognostically similar to SCC; clinicopathologic correlation is recommended in these cases.

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.

診斷 初步評估 分期 初始治療 療效評估 輔助治療 追蹤



§ : Image studies is indicated for extensive disease (deep structural involvement such as bone, deep soft tissue, perineural disease)

+ : regional or distal metastatic disease(初始皮膚病灶治療同侷限型)

☆ : 附件一

△ : 附件二


: T any, N0, M0(附件三)



皮膚癌
多專科團隊

基底細胞癌(BCC)

復發



Hedgehog pathway inhibitor : Vismodegib/ Sonidegib
 Programmed cell death protein 1 (PD-1) inhibitor (cemiplimab)





FOLLOW-UP

- H&P
 - ▶ Including complete skin exam every 6–12 mo for the first 5 years, and then at least annually for life^u
- Consider imaging if clinical exam is insufficient for following the disease^v
- Patient education:
 - ▶ Sun protection
 - ▶ Self-examination

RECURRENCE OR ADVANCED DISEASE

Local recurrence

Follow Primary Treatment pathway for high-risk disease (BCC-3)

Primary or recurrent nodal metastases

Multidisciplinary consultation to consider one or more of the following options:
Surgery
or
If surgery is not feasible then RT^j or systemic therapy^q

- Hedgehog pathway inhibitor (HHI)
 - ▶ Vismodegib
 - ▶ Sonidegib (category 2B)
- Programmed cell death protein 1 (PD-1) inhibitor (cemiplimab-rwlc)^w
- Clinical trial

Distant metastases

Multidisciplinary consultation to consider:
Systemic therapy^q

- HHI
 - ▶ Vismodegib
- PD-1 inhibitor (cemiplimab-rwlc)^w

or
RT^j or surgery for limited metastatic disease^x
or
Palliation and best supportive care

^j Principles of Radiation Therapy (BCC-D).

^q Principles of Systemic Therapy (BCC-E).

^u Follow-up with a dermatologist is strongly recommended if any of the following criteria are met: past or imminent solid organ, marrow, or stem cell transplant, one or more cutaneous melanomas in the past 5 years, or four or more non-melanoma skin cancers in the past 5 years.

^v Imaging modality and targeted area should be at the discretion of the treating team based on the suspected extent of disease (ie, local, regional, metastatic). Histologic confirmation is often sufficient to diagnose local recurrence, but MRI can be considered to assess extent of local disease. For nodal or distant metastasis, histologic analysis and/or CT imaging can be used for confirmation and to gauge extent of disease.


^w Cemiplimab-rwlc is recommended for patients with locally advanced or metastatic basal cell carcinoma (mBCC) previously treated with an HHI or for whom an HHI is not appropriate.

^x Under highly selective circumstances, in the context of multidisciplinary consultation, resection of limited metastases can be considered.

Note: All recommendations are category 2A unless otherwise indicated.
Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



皮膚癌
多專科團隊



^a [Principles of Pathology \(BCC-A\)](#).

^b [Risk Factors for Recurrence \(BCC-B\)](#).

^e Any high-risk factor places the patient in the high-risk category.

^h [Principles of Treatment \(BCC-C\)](#).

^j [Principles of Radiation Therapy \(BCC-D\)](#).

^k Determination of the appropriateness of RT should be performed by a radiation oncologist.

ⁿ Mohs surgery should be performed by dermatologic surgeons who have specialized training and experience in this procedure.

^o As per other appropriate use criteria. Task Force/Committee Members, Vidal CI, et al. J Am Acad Dermatol 2019;80:189-207.

^p PDEMA with permanent section analysis or intraoperative frozen section analysis is an alternative to Mohs. See Principles of PDEMA Technique (SCC-G) within the [NCCN Guidelines for Squamous Cell Skin Cancer](#).

^q For clinically diagnosed non-facial BCCs <6 mm in depth on the head, neck, hands, feet, pretibial, and anogenital area that are clinically confined to the dermis, C&E or shave removal may be considered as an alternative primary treatment option if Mohs, resection with PDEMA, and standard excision are difficult to perform due to patient comorbidities (eg, thrombocytopenia, immunosuppression, bleeding diathesis, multiple primary BCCs). See [Risk Factors for Recurrence \(BCC-B\)](#).

^v Fraga SD, et al. Dermatol Surg 2022;48:704-710.

^w Under highly selective circumstances, in the context of multidisciplinary consultation, resection of limited metastases can be considered.


Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



FOLLOW-UP

RECURRENCE



^x Follow-up with a dermatologist is strongly recommended if any of the following criteria are met: past or imminent solid organ, marrow, or hematopoietic cell transplant; one or more cutaneous melanomas in the past 5 years; or four or more non-melanoma skin cancers in the past 5 years.

^y Imaging modality and targeted area should be at the discretion of the treating team based on the suspected extent of disease (ie, local, regional, metastatic). Histologic confirmation is sufficient to diagnose local recurrence, but MRI can be considered to assess extent of local disease. For nodal or distant metastasis, histologic analysis and/or CT imaging can be used for confirmation and to gauge extent of disease.

Note: All recommendations are category 2A unless otherwise indicated.
Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.

基底細胞癌(BCC)

癌症藥物停藥準則

- 根據CTCAE (Common Terminology Criteria for Adverse Events, Version 4.0 Published: May 28, 2009 【v4.03: June 14, 2010】)，出現Grade 3 ~ Grade 4 adverse event。
- 停藥至adverse event回復至Grade 1或Baseline時可再次用藥，但有些患者必須調整用藥劑量。
- 特定藥物治療下疾病仍持續進展，根據追蹤及評估顯示疾病對此特定藥物治療無效（考慮停止投藥並選擇其他治療方法）。
- 病患要求 (Hospice care或其他因素)。
- 病患死亡。



基底細胞癌(BCC)

附件一：

STRATIFICATION TO DETERMINE TREATMENT OPTIONS FOR LOCAL BCC BASED ON RISK FACTORS FOR RECURRENCE¹

Risk Group	Low Risk	High Risk
Treatment Options	See BCC-2	See BCC-3
H&P		
Location/size	Trunk, extremities <2 cm	Trunk, extremities ≥2 cm Cheeks, forehead, scalp, neck, and pretibia (any size) Head, neck, hands, feet, pretibia, and anogenital (any size) ³
Borders	Well-defined	Poorly defined
Primary vs. recurrent	Primary	Recurrent
Immunosuppression	(-)	(+)
Site of prior RT	(-)	(+)
Pathology (See BCC-A)		
Subtype	Nodular, superficial ²	Aggressive growth pattern ⁴
Perineural involvement	(-)	(+)

¹ Any high-risk factor places the patient in the high-risk category.
² Low-risk histologic subtypes include nodular, superficial, and other non-aggressive growth patterns such as keratotic, infundibulocystic, and fibroepithelioma of Pinkus.
³ This area constitutes high risk based on location, independent of size. Narrow excision margins due to anatomic and functional constraints are associated with increased recurrence rates with standard histologic processing. Complete margin assessment such as with Mohs or PDEMA is recommended for optimal tumor clearance and maximal tissue conservation. For tumors <6 mm in size, without other high-risk features, other treatment modalities may be considered if at least 4-mm clinically tumor-free margins can be obtained without significant anatomic or functional distortions.
⁴ Having (mixed) infiltrative, micronodular, morpheaform, basosquamous, sclerosing, or carcinosarcomatous differentiation features in any portion of the tumor. In some cases basosquamous tumors may be prognostically similar to SCC; clinicopathologic correlation is recommended in these cases.

Note: All recommendations are category 2A unless otherwise indicated.
Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



基底細胞癌(BCC)

附件二:



National Comprehensive Cancer Network®

NCCN Guidelines Version 1.2022 Basal Cell Skin Cancer

[NCCN Guidelines Index](#)
[Table of Contents](#)
[Discussion](#)

PRINCIPLES OF RADIATION THERAPY

General Principles

- Protracted fractionation is associated with improved cosmetic results and should be utilized for poorly vascularized or cartilaginous areas.
- RT is contraindicated for genetic conditions predisposing to skin cancer (eg, basal cell nevus syndrome) and relatively contraindicated for patients with connective tissue diseases (eg, scleroderma).
- Given higher complication rates, re-irradiation should not be routinely utilized for recurrent disease within a prior radiation field.
- Isotope-based brachytherapy can be an effective treatment for certain sites of disease, particularly on the head and neck.
- There are insufficient long-term efficacy and safety data to support the routine use of electronic surface brachytherapy.

General Treatment Information

Dosing Prescription Regimen

<u>Definitive RT</u>	<u>Examples of Electron Beam Dose and Fractionation</u>
Tumor diameter <2 cm	60–64 Gy over 6–7 weeks 50–55 Gy over 3–4 weeks 40 Gy over 2 weeks 30 Gy in 5 fractions over 2–3 weeks
Tumor diameter ≥2 cm, T3/T4, or those with invasion of bone or deep tissue	60–70 Gy over 6–7 weeks 45–55 Gy over 3–4 weeks
<u>Postoperative Adjuvant RT</u>	60–64 Gy over 6–7 weeks 50 Gy over 4 weeks

Note: All recommendations are category 2A unless otherwise indicated.
Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.

Version 1.2022, 11/17/21 © 2021 National Comprehensive Cancer Network® (NCCN®). All rights reserved. NCCN Guidelines® and this illustration may not be reproduced in any form without the express written permission of NCCN.

BCC-D

皮膚癌
多專科團隊

基底細胞癌(BCC)

附件三-1:

現在使用AJCC第八版的Skin tumor在頭頸部及生殖部位才需要staging

Staging

Table 1
 American Joint Committee on Cancer (AJCC)
TNM Staging Classification for Cutaneous Squamous Cell Carcinoma (cSCC) and Other Cutaneous Carcinomas
 (7th ed., 2010)
Primary Tumor (T)*
TX Primary tumor cannot be assessed
T0 No evidence of primary tumor
Tis Carcinoma in situ
T1 Tumor 2 cm or less in greatest dimension with less than two high-risk features**
T2 Tumor greater than 2 cm in greatest dimension or
 Tumor any size with two or more high-risk feature
T3 Tumor with invasion of maxilla, mandible, orbit, or temporal bone
T4 Tumor with invasion of skeleton (axial or appendicular) or perineural invasion of skull base
 *Excludes cSCC of the eyelid
 ** High-risk features for the primary tumor (T) staging
 Depth/invasion > 2 mm thickness
 Clark level ≥ IV
 Perineural invasion
 Anatomic location Primary site ear
 Primary site non-hair-bearing lip
 Differentiation Poorly differentiated or undifferentiated

Regional Lymph Nodes (N)
NX Regional lymph nodes cannot be assessed
N0 No regional lymph node metastases
N1 Metastasis in a single ipsilateral lymph node, 3 cm or less in greatest dimension
N2 Metastasis in a single ipsilateral lymph node, more than 3 cm but not more than 6 cm in greatest dimension; or in multiple ipsilateral lymph nodes, none more than 6 cm in greatest dimension; or in bilateral or contralateral lymph nodes, none more than 6 cm in greatest dimension
N2a Metastasis in a single ipsilateral lymph node, more than 3 cm but not more than 6 cm in greatest dimension
N2b Metastasis in multiple ipsilateral lymph nodes, none more than 6 cm in greatest dimension
N2c Metastasis in bilateral or contralateral lymph nodes, none more than 6 cm in greatest dimension
N3 Metastasis in a lymph node, more than 6 cm in greatest dimension

Distant Metastasis (M)
M0 No distant metastases
M1 Distant metastases

[Continue](#)

Used with the permission of the American Joint Committee on Cancer (AJCC), Chicago, Illinois. The original and primary source for this information is the AJCC Cancer Staging Manual, Seventh Edition (2010) published by Springer Science and Business Media LLC (SBM). (For complete information and data supporting the staging tables, visit www.springer.com.) Any citation or quotation of this material must be credited to the AJCC as its primary source. The inclusion of this information herein does not authorize any reuse or further distribution without the expressed, written permission of Springer SBM, on behalf of the AJCC.

基底細胞癌(BCC)

附件三-2:



NCCN Guidelines Version 1.2014 Basal and Squamous Cell Skin Cancers

[NCCN Guidelines Index](#)
[Basal and Squamous Cell TOC](#)
[Discussion](#)

Table 1 Continued
American Joint Committee on Cancer (AJCC)
TNM Staging Classification for Cutaneous Squamous Cell Carcinoma (cSCC) and Other Cutaneous Carcinomas (7th ed., 2010)

Anatomic Stage/Prognostic Groups

Stage 0	Tis	N0	M0
Stage I	T1	N0	M0
Stage II	T2	N0	M0
Stage III	T3	N0	M0
	T1	N1	M0
	T2	N1	M0
Stage IV	T3	N1	M0
	T1	N2	M0
	T2	N2	M0
	T3	N2	M0
	T Any	N3	M0
	T4	N Any	M0
	T Any	N Any	M1

Histologic Grade (G)

GX	Grade cannot be assessed
G1	Well differentiated
G2	Moderately differentiated
G3	Poorly differentiated
G4	Undifferentiated

Used with the permission of the American Joint Committee on Cancer (AJCC), Chicago, Illinois. The original and primary source for this information is the AJCC Cancer Staging Manual, Seventh Edition (2010) published by Springer Science and Business Media LLC (SBM). (For complete information and data supporting the staging tables, visit www.springer.com.) Any citation or quotation of this material must be credited to the AJCC as its primary source. The inclusion of this information herein does not authorize any reuse or further distribution without the expressed, written permission of Springer SBM, on behalf of the AJCC.

Reference

1. NCCN Clinical Practice Guideline in Oncology, Basal and Squamous Cell Skin Cancers, Version 1.2022.
2. G Ital Dermatol Venereol. 2016 Feb;151(1):77-86. Epub 2014 Jun 30. Treatments of advanced basal cell carcinoma: a review of the literature.
3. Sekulic A, Migden MR, Basset-Seguín N, et al. Long-term safety and efficacy of vismodegib in patients with advanced basal cell carcinoma (aBCC): 18-month update of the pivotal ERIVANCE BCC study. ASCO Meeting Abstracts 2013;31:9037.
4. Mendenhall WM, Ferlito A, Takes RP, et al. Cutaneous head and neck basal and squamous cell carcinomas with perineural invasion. Oral Oncol 2012;48:918-922.
5. Sekulic A, Migden MR, Oro AE, et al. Efficacy and safety of vismodegib in advanced basal-cell carcinoma. N Engl J Med 2012;366:2171-2179.
6. Lott DG, Manz R, Koch C, Lorenz RR. Aggressive behavior of nonmelanotic skin cancers in solid organ transplant recipients. Transplantation 2010;90:683-687.
7. Basosquamous carcinoma. J Am Acad Dermatol 2009;60:137-143.
8. Mendenhall WM, Amdur RJ, Hinerman RW, et al. Radiotherapy for cutaneous squamous and basal cell carcinomas of the head and neck. Laryngoscope 2009;119:1994-1999.
9. Mosterd K, Krekels GA, Nieman FH, et al. Surgical excision versus Mohs' micrographic surgery for primary and recurrent basal-cell carcinoma of the face: a prospective randomised controlled trial with 5-years' follow-up. Lancet Oncol 2008;9:1149-1156.
10. Neville JA, Welch E, Leffell DJ. Management of nonmelanoma skin cancer in 2007. Nat Clin Pract Oncol 2007;4:462-469.
11. Rodriguez-Vigil T, Vazquez-Lopez F, Perez-Oliva N. Recurrence rates of primary basal cell carcinoma in facial risk areas treated with curettage and electrodesiccation. J Am Acad Dermatol 2007;56:91-95.
12. Rubin AI, Chen EH, Ratner D. Basal-cell carcinoma. N Engl J Med 2005;353:2262-2269.
13. Bath-Hextall F, Bong J, Perkins W, Williams H. Interventions for basal cell carcinoma of the skin: systematic review. BMJ 2004;329:705.